

Landslide Atlas of India

भारत के भूस्खलन एटलस

(Mapping, Monitoring and R&D studies using Remote Sensing data)

(सुदूर संवेदन आधारित मानचित्रण, निगरानी और अनुसंधान एवं विकास)



राष्ट्रीय सुदूर संवेदन केंद्र / National Remote Sensing Centre
भारतीय अंतरिक्ष अनुसंधान संगठन / Indian Space Research Organisation
अंतरिक्ष विभाग / Department of Space
हैदराबाद / Hyderabad-500037



nrsc

February 2023

Landslide Atlas of India

भारत के भूस्खलन एटलस

(Mapping, Monitoring and R&D studies using Remote Sensing data)

(सुदूर संवेदन आधारित मानचित्रण, निगरानी और अनुसंधान एवं विकास)



राष्ट्रीय सुदूर संवेदन केंद्र / National Remote Sensing Centre
भारतीय अंतरिक्ष अनुसंधान संगठन / Indian Space Research Organisation

अंतरिक्ष विभाग / Department of Space

हैदराबाद / Hyderabad-500037

February 2023



nrsc



Copyright © 2023

National Remote Sensing Centre, ISRO

Published by

National Remote Sensing Centre, ISRO
Department of Space, Government of India
Hyderabad – 500 037, Telangana, India
<https://www.nrsc.gov.in>

Printed By

National Remote Sensing Centre, ISRO Hyderabad, India.

Photos

Front Cover page: High-resolution Image of landslide crown area of the Puthumala landslide taken from UAV.

Back Cover Page: Field photographs of landslides from different parts of Uttarakhand.

Production

NRSC has played a key role in the DMS programme of ISRO in providing space-based support to various landslide disasters since 2005 in a programmatic manner. NRSC has prepared landslide hazard zonation (LHZ) map and Atlas for the first time in India on 1:25000 scale in the year 2001. Damage assessment for major landslide disasters was carried out using satellite data and aerial images and value-added products are provided to stake holders. NRSC has completed Seasonal Landslide Inventory Mapping Project (SLIM) for India for the post-monsoon of 2014 and 2017.

Reproduction

The publication may be produced in whole or in part and in any form of educational or non-profit purposes without permission from the copyright holder, provided the acknowledgement and citation (mentioned below) of the source is made in all type of publication, which include research articles, journal papers, conference proceeding, books, book chapter, scientific reports, dissertation, etc. NRSC would appreciate receiving a copy of any publication that uses this publication as a source. No use of this publication may be made for resale or for any other commercial purposes. The view and interpretation in this publication are those of the author(s). They are not attributable to NRSC and do not imply the expression of any opinion concerning the legal status of any country, territory, city or area of its authorities or concerning the delimitation of its frontiers or boundaries or the endorsement of any product. The administrative boundaries presented in this publication are only for scientific study and not for statutory purpose.

Citation

Nirmala Jain, Priyom Roy, Tapas Ranjan Martha, Punit Jalan and Aishwarya Nanda (2023). Landslide Atlas of India (Mapping, Monitoring and R&D studies using Remote Sensing data). NRSC special publication. NRSC/ISRO. Document number: NRSC-RSA-GSG-GMED-FEB 2023-TR-0002167-V1.0

Printed at Printing Facility, PPEG, NRSC

DOCUMENT CONTROL SHEET

1	Security Classification	Official			
2	Distribution	General			
3	Report / Document version	(a) Issue no. 01	(b) Revision & Date	27.02.2023	
4	Report / Document Type	Technical report			
5	Document Control Number	NRSC-RSA-GSG-GMED-FEB 2023-TR-0002167-V1.0			
6	Title	Landslide atlas of India (Mapping, Monitoring and R&D studies using Remote Sensing data)			
7	Particulars of collation	Pages 76	Figures 55	Tables 04	References 16
8	Author(s)	Nirmala Jain, Priyom Roy , Tapas Ranjan Martha, Punit Jalan and Aishwarya Nanda			
9	Affiliation of authors	Geohazards and Mineral Exploration Division, Geosciences Group, RSAA, NRSC/ISRO			
10	Scrutiny mechanism	Compiled by Nirmala Jain S/E 'SE' Priyom Roy S/E 'SE'	Reviewed by Dr. I. C. Das Group Head, GSG	Approved and Controlled Dr. K Vinod Kumar Deputy Director (RSAA)	
11	Originating unit	National Remote Sensing Centre, Hyderabad-500037			
12	Sponsor(s)/ Name and Address	Indian Space Research Organisation			
13	Date of Initiation	01 November 2022			
14	Date of Publication	27 February 2023			
15		<p>Abstract: This Atlas provides the details of landslide present in Landslide provinces of India including damage assessment of specific landslide locations. Geospatial landslide inventory database consisting of ~80,000 landslides in India mapped by NRSC/ISRO under its DMS programme during the 1998-2022 period is reported here. The database covers landslide vulnerable regions in 17 states and 02 UTs of India in the Himalayas and Western Ghats. The database includes three types of landslide inventory – seasonal, event-based and route-wise for the 1998-2022 period. Seasonal inventory contains pan-India landslide database corresponding to the 2014 and 2017 rainy season in India. Event-based inventory contains details of some of the major triggering events such as Kedarnath and Kerala disasters, and Sikkim earthquake as well as few large valley blocking landslides. Route-wise inventory contains details of landslides along selected routes of tourist and pilgrimage importance. Satellite data of high to very high resolution such as IRS-1D PAN+LISS-III, Resourcesat-1, 2 and 2A LISS-IV Mx, Cartosat-1 and 2S, data from International satellites (Sentinel-1&2, Pleiades and WorldView) and Aerial images were used in the mapping of landslides. Some of the mapped landslides were validated in the field using mobile App and news reports. The database were used to rank 147 districts in 17 states and 02 UTs of India for their exposure to landslides in terms of key socio-economic parameters. Lastly advanced techniques in landslide detection, modeling and prediction are also explained.</p> <p>Keywords: Landslides, Remote Sensing, Landslide Provinces in India, Landslide Inventory</p>			

भारतीय अन्तरिक्ष अनुसंधान संगठन
अन्तरिक्ष विभाग
भारत सरकार
अन्तरिक्ष भवन
न्यू बी ई एल रोड, बेंगलूर - 560 231, भारत
दूरभाष : +91-80-2341 5241 / 2217 2333
फैक्स : +91-80-2341 5328



Indian Space Research Organisation
Department of Space
Government of India
Antariksh Bhavan
New BEL Road, Bangalore - 560 231, India
Telephone: +91-80-2341 5241 / 2217 2333
Fax : +91-80-2341 5328
e-mail : chairman@isro.gov.in

Shri S. Somanath
Chairman ISRO, Secretary DOS



Message

India is prone to many natural disasters like flood, landslide, cyclone, forest fire, earthquake, drought etc. Satellites provide synoptic observations of natural disasters at regular intervals that help in management of disasters. In order to better understand the risks due to such disasters, it is also necessary to integrate satellite and field-based observations and to work towards risk reduction principles. Satellite communication and navigation systems also play an important role in disaster management with improved technological options.

Disaster Management Support (DMS) Programme of ISRO, comprehensively addresses various aspects of natural disasters in the country, using space-based inputs. ISRO disseminates relevant information in the interactive geospatial domain through various portals like Bhuvan, National Database for Emergency Management and MOSDAC for the administrators to better understand the impact and for improved decision support.

ISRO provides satellite-based near real-time information support to Central Ministries / Departments and State Ministries / Departments, prior to and after major natural disasters. Under the DMS programme, ISRO has been actively working in landslide disaster response and mitigation. Satellite imagery and damage analysis have been provided for major landslide disasters in recent times in Kedarnath, Kerala, Chamoli and Assam.

The "Landslide Atlas of India" created by National Remote Sensing Centre, Hyderabad is another value addition in the field of landslide response and research. It illustrates the overall landslide scenario in India and also puts forward key research outcomes in the field of landslide hazard. I congratulate team NRSC for this effort and hope this atlas will be beneficial to disaster responders and researchers.

Bengaluru
Feb 27, 2023

(S. Somanath)



nrsc

Dr. Prakash Chauhan
Director, NRSC



FOREWORD

NRSC has played a pivotal role in the DMS programme of ISRO in providing space-based support to various natural disasters since 2005 in a programmatic manner. Many projects, both operational and R&D nature, are executed by NRSC as a part of the ISRO's DMS programme.

In the field of landslide studies, NRSC has prepared a landslide hazard zonation (LHZ) map on 1:25,000 scale in the year 2001 for the first time in India using GIS and AHP techniques. Damage assessment for major landslide disasters was carried out using satellite data and aerial images and value-added products are provided to stakeholders. NRSC has completed Seasonal Landslide Inventory Mapping Project (SLIM) for India for the post-monsoon season of 2014 and 2017. Pan-India rainfall-triggered landslide databases are available for the first time in the country as a result of this project. NDMA has assigned the responsibility of rapid landslide mapping using satellite data to NRSC through office memorandum no. 6-66/2013-Mit-II/4531, dated 18 Jan 2017. New research activities were taken up by NRSC, particularly, understanding the kinematics of landslides using PSInSAR technique that will be a new paradigm for landslide early warning in near future.

The "Landslide Atlas of India" created by NRSC showcases the recent responses and developments in the landslide disaster in the country using space data and geospatial models. The state-wise landslide maps show the geospatial distribution of landslides. Few representative large landslides from the Himalaya and Western ghats have been showcased using high-resolution satellite imagery. I congratulate the team for compiling this atlas and hope it will prove to be beneficial to researchers and experts of landslides.

Hyderabad
Feb 27, 2023

(Prakash Chauhan)



nrsc

Dr. K.Vinod Kumar
Dy. Director
Remote Sensing Applications (RSA)
NRSC



PREFACE

Landslides are controlled and triggered by geological, topographical and hydrological factors. Studying landslide hazard using satellite data has an inherent advantage since it occurs mostly in remote and inaccessible mountains. NRSC has developed expertise in addressing various aspects of landslide hazard using both optical and SAR satellite images.

NRSC has created a pan India database of ~80,000 landslides those occurred during last two decades using satellite data. This database contains both seasonal landslides and event-based landslides. These data were created as polygons in GIS, and contains several attribute information relevant for management of landslide disaster. This pan India database of landslides helped to understand the critical lithology, geological structure and rainfall responsible for occurrence of landslides. This database also helped to understand the district-wise landslide exposure in India. Other than the landslide database, various technological advancements such as debris flow modeling using UAV, landslide volume estimation, landslide mobile app and landslide kinematics using InSAR were also developed by NRSC.

The “Landslide Atlas of India” prepared by NRSC is a rich collection of landslide inventory which will help user departments in prioritisation of developmental activities in tectonically and ecologically sensitive mountainous areas in India. I congratulate the team for preparing this atlas.

Hyderabad
Feb 27, 2023

(K Vinod Kumar)

ACKNOWLEDGEMENTS

The “Landslide Atlas of India” is a consolidation of work carried out by NRSC related to landslide disasters under the Disaster Management Support (DMS) programme of ISRO. We sincerely thank Shri S Somanath, Secretary DOS and Chairman ISRO, Dr. Prakash Chauhan, Director NRSC for their encouragement and support to this work. We thank Dr. Shantanu Bhatawdekar, Scientific Secretary ISRO and Director EDPO, Dr. K Vinod Kumar, DD (RSA) NRSC, Dr. John Mathew, Associate Director EDPO and Dr. K H V Durga Rao, Group Director, DMSG NRSC for their active support to this study.

Satellite data are the main requirement for this work carried out by NRSC. Hence, we thank Dr. N Aparna, Group Head NDC, NRSC and her team for providing the IRS satellite data. We also thank Copernicus, USGS, International Charter Space and Major Disasters for providing satellite data for emergency response to landslide disaster events in India. We also thank Shri M. Arulraj and Bhuvan team for their support in the web-based cataloguing of landslides. We thank our colleagues in Geosciences Group, NRSC and knowledge partner institutions (GSI, NDMA, SDMAs, NIDM, CBRI, WIHG, NRSC-RRSCs, NRSC-ASDMA, IIRS, SAC, NESAC, CRRI, DGRE, IITs, ITC-Netherlands, State remote sensing centres etc.) in India who have contributed to this work.

Dr. Iswar Chandra Das
Group Head, Geosciences Group

Table of Contents

1. Message	i
2. Foreword	iii
3. Preface	v
4. Acknowledgements	vii
5. List of Figures	1
6. List of Tables	5
7. Introduction and Background	7
8. Landslide Disaster in India	8
9. Landslide Classification	9
10. Rapid Response to Landslide Disaster using EO Data	11
11. Landslide Inventory of India	17
12. Landslide Susceptibility Mapping and Districts-wise Landslide Risk Exposure	55
13. Research & Development Studies in Landslides Studies	63
14. References	71
15. Bibliography of landslide studies done by NRSC	73

List of figures

Figure 1. Block diagram of features of landslide: earth flow type.

Figure 2. Landslides from different parts of India.

Figure 3. Landslide damage to New Haflong Railway Station Dima Hasao District of Assam (15-16 May 2022).

Figure 4. Damage caused by the Raunthi Gad landslide. Synoptic view of the affected region as seen from Resourcesat-2A LISS IV image (08 February 2021) shown in FCC.

Figure 5. Damage caused by the Raunthi Gad landslide. Field observations.

Figure 6. Geomorphological changes leading to the disaster around Kedarnath, Uttarakhand (2013).

Figure 7. Blockage of road (highlighted with circles) adjacent to the Tista river due to landslides triggered by Sikkim Earthquake, 2011.

Figure 8. Parts of the road destroyed by landslides triggered due to the earthquake near Chungthang area in north Sikkim.

Figure 9. Landslide damage in Okhimath town. GeoEye-1 images showing buildings that were damaged (yellow outlines) by landslides.

Figure 10. Landslides inventory of India.

Figure 11. Landslides mapped using high-resolution satellite data in Jammu & Kashmir and Ladakh, which occurred between 2014 to 2017.

Figure 12. Paryote landslides near Paryote village in Doda district, Jammu and Kashmir.

Figure 13. Landslides mapped using high-resolution satellite data in Himachal Pradesh which were occurred between 1998 to 2017.

Figure 14. Kotropi landslides, Mandi district, Himachal Pradesh.

Figure 15. Landslides mapped using high-resolution satellite data in Uttarakhand, which occurred between 1998 to 2022.

Figure 16. Landslides around Kedarnath, Uttarakhand.

Figure 17. Landslides mapped using high-resolution satellite data in Haryana, which occurred in 2017.

Figure 18. Panchkula landslides, Haryana.

Figure 19. Landslides mapped using high-resolution satellite data in Sikkim, which occurred between 2011 to 2022.

Figure 20. Mantam landslide, Sikkim.

Figure 21. Landslides mapped using high-resolution satellite data in West Bengal, which occurred between 2011 to 2017.

Figure 22. Landslide near Kurseong village, West Bengal.

Figure 23. Landslides mapped using high-resolution satellite data in Arunachal Pradesh, which occurred between 2014 to 2021.

Figure 24. Landslide from West Siang, Arunachal Pradesh.

Figure 25. Landslides mapped using high-resolution satellite data in Assam, which occurred between 2014 to 2022.

Figure 26. Landslide from Dima Hasao District, Assam.

Figure 27. Landslides mapped using high-resolution satellite data in Nagaland, which occurred between 2014 to 2018.

Figure 28. Landslide from Kikrumba in Phek district, Nagaland.

Figure 29. Landslides mapped using high-resolution satellite data in Manipur, which occurred between 2014 to 2022.

Figure 30. Landslide near Tupul Railway Station, Noney, Manipur.

Figure 31. Landslides mapped using high-resolution satellite data in Mizoram, which occurred between 2014 to 2018.

Figure 32. Landslide from Aizwal, Mizoram.

Figure 33. Landslides mapped using high-resolution satellite data in Tripura, which occurred between 2014 to 2018.

Figure 34. Landslide from Dhalai, Tripura.

Figure 35. Landslides mapped using high-resolution satellite data in Meghalaya, which occurred between 2014 to 2017.

Figure 36. Landslide from East Khasi Hills, Meghalaya.

Figure 37. Landslides mapped using high-resolution satellite data in Maharashtra, which occurred between 2014 to 2021.

Figure 38. Landslide from Taliye Village, Mahad district, Maharashtra.

Figure 39. Landslides mapped using high-resolution satellite data in Karnataka, which occurred between 2014 to 2018.

Figure 40. Landslide from Kodagu and Dakshina Kannada districts, Karnataka.

Figure 41. Landslides mapped using high-resolution satellite data in Kerala, which occurred between 2014 to 2021.

Figure 42. Landslide from Puthumala, Wayanad district, Kerala.

Figure 43. Landslides mapped using high-resolution satellite data in Tamil Nadu, which occurred between 2014 to 2018.

Figure 44. Landslides in Coimbatore district, Tamil Nadu.

Figure 45. Landslides mapped using high-resolution satellite data in Goa, which occurred between 2014 to 2017.

Figure 46. LSZ map for Shimla– Rampur– Sarahan – Sumdo Route.

Figure 47. Socio-economic parameter risk exposure map, The contribution of each exposure element for the top ten landslide-exposed districts is shown in bar diagram.

Figure 48. Landslide prediction using MT InSAR. Kikruma landslide.

Figure 49. Flow modeling for Puthumala Landslide, Wayanad District, Kerala.

Figure 50. Landslide modeling using UAV data, Puthumala Landslide, Wayanad District, Kerala.

Figure 51. Landslide susceptibility maps generated using spatial sampling strategy. Landslide susceptibility maps of Mizoram.

Figure 52. Landslide early warning for advance 3 days given in parts of Himachal Pradesh.

Figure 53. Step-by-step procedure for landslide data collection using FLIM mobile app.

Figure 54. Landslide data from western Maharashtra collected in Bhuvan FLIM database.

Figure 55. Ground landslide data collected in NER using FLIM mobile App.

List of tables

Tables

Table 1. The classification of landslides by Varnes (1978).

Table 2. Landslides inventory of India statistic.

Table 3. Table shows the rank of districts in 17 states of India for their exposure to landslides .

Table 4. Table showing triggering probability for landslide trigger.

Introduction and background

The frequently used definition of landslide is “a movement of mass of rock, earth or debris down a slope” (Cruden, 1991). Different types of landslides such as Debris slide, debris flows, Rock slide, rock fall etc. can be triggered by rainfall, undercutting of slopes due to flooding or excavation, earthquakes, snowmelt and other natural causes, as well as anthropogenic causes such as over grazing by cattle, terrain cutting and filling, excessive development, etc. They can occur on many types of terrain given the right conditions of soil, rock, geological structure and slope.

Landslides are among the main natural catastrophes, which cause major problems in mountainous terrain by killing hundreds of people every year besides damaging property, disrupting transportation and blocking communication links. In some areas, such as the western coastal parts of North and South America, Central America, Alpine regions of Italy, France, Switzerland and Austria in Europe, Himalayan regions of India, Nepal in Asia and parts Central Asia, the effects of landslides are more pronounced mainly due to the spurred developmental activities to meet the ever growing demand of people. As per the official figures of United Nations International Strategy for Disaster Reduction (UN/ISDR) and the Centre for Research on the Epidemiology of Disasters (CRED) for the year 2006, landslide ranked 3rd in terms of number of deaths among the top ten natural disasters. Approximately 4 million people were affected by landslides in 2006 (OFDA/CRED, 2006). Regions with the highest landslide risk can be found in Colombia, Tajikistan, India, and Nepal where the estimated number of people killed per year per 100 km² was found to be greater than one (Nadim et al., 2006)

India is among the top four countries with highest landslide risk, where the estimated loss of life per 100 km² per year is greater than one. In India, most of the landslides occur in the monsoon months (between June and September). Globally, India contribute 16 % rainfall-triggered landslide events (Source: Froude and Petley, 2018). Landslides mainly occur along slope of hills due to natural triggering factors such as heavy rainfall and earthquake or due to anthropogenic activities such as construction of roads, buildings and mining. The atlas comprises of information about the landslide records occurred between 1998 to 2022. It highlights the understanding of socio-economic parameters with respect to district wise landslide distribution in the country.

Landslide disaster in India

Landslides are one of the common geological hazards in hilly areas throughout the world. Other than geological and anthropogenic causes, rainfall is the natural triggering factor for occurrence of landslides. Rainfall-induced landslides are the result of the combined action of water on topography, geology, soil and vegetation. Landslides have led to massive environmental damages such as increase in sediment discharge due to soil erosion and loss of human lives every year. India, a country with varied physiographic and climatic conditions, frequently faces the vagaries of landslide disaster. Approximately 0.42 million sq. km or 12.6% of land area, excluding snow covered area, is prone to landslide hazard. Out of this, 0.18 million sq. km falls in North East Himalaya, including Darjeeling and Sikkim Himalaya; 0.14 million sq. km falls in North West Himalaya (Uttarakhand, Himachal Pradesh and Jammu & Kashmir); 0.09 million sq. km in Western Ghats and Konkan hills (Tamil Nadu, Kerala, Karnataka, Goa and Maharashtra) and 0.01 million sq. km in Eastern Ghats of Aruku area in Andhra Pradesh (www.gsi.gov.in). In India, landslides mostly occur in the monsoon season. Himalayas and Western Ghats are highly susceptible to mass movements due to hilly topography and heavy rainfall

The majority of landslides are triggered by variability in rainfall patterns, while sporadic events such as very heavy rainfall outside the monsoon period (Kedarnath event of 2013) and earthquakes (Sikkim earthquake) cause significant disruption to livelihood and infrastructure. The Northernmost Indian states, J&K, Himachal Pradesh, and Uttarakhand, are the worst affected states by the landslides disasters as the majority of the territory falls within the Himalayas. Many districts of these states have high population densities, and major pilgrimage routes or major tourism spots are exposed and affected by landslides. Although the North Eastern states have many landslides yearly, they are not particularly vulnerable to them in terms of socioeconomic factors due to their lower population density and wide unoccupied mountain areas.

The vulnerability of the inhabitants and households is more significant in the Western Ghats due to the very high population and household density, especially in Kerala, even when fewer landslides exist than in the Himalayan regions. The Himalayan range has a significant influence on the geographic distribution of landslides. The spatial distribution of landslides is controlled by the shale-sandstone-metasediments generating dissected hills and valleys over the Himalayan range with slopes 30° to 45°. This, together with rainfall between 750-1000 mm is the cause of landslides. Also, the Chenab subbasin, whose geographic expanse primarily comprises steeply sloping mountains, records the highest number of landslides. The steep escarpments of the Western Ghats record occurrences of landslides, but however, these are primarily controlled by the soil cover on the slopes.

Landslide Classification

Table 1. The classification of landslides.

Abbreviated Classification of Slope Movements				
Type of movement		TYPE OF MATERIAL		
		Bed Rock	Engineering Soils	
			Predominantly Course	Predominantly Fine
Fall		Rock Fall	Debris Fall	Earth Fall
Topple		Rock Topple	Debris Topple	Earth Topple
Slide	Rotational (Slump)		Rotational Debris Slide	Rotational Earth Slide
	Translational	Translational Rock Slide	Translational Debris Slide	Translational Earth Slide
Lateral Spread		Rock Spread	Debris Spread	Earth Spread
Flow		Rock Flow (Deep creep)	Debris Flow (Soil Creep)	Earth Flow (Soil Creep)
Complex	Combination of two or more principal types of movement			




Figure 1. Block diagram of features of landslide: Earth flow type (pubs.usgs.gov).

Landslide Classification

Types of Movement

Falls: Falls are rapid movements of rocks and boulders detached from steep slopes or cliffs along fractures, joints, and bedding planes.

Topple: It is the forward rotation of a mass of debris or rock out of a slope. The slope failure generally occurs at point near the base of the block of rock.

Slides: A slide is a downslope movement of material that occurs along a slip surface.

Rotational slide: In this slide the slide movement is roughly rotational about an axis which is parallel to the ground surface and transverse across the slide.

Translational slide: In this slide the landslide mass moves along a planar surface with rotation or backward tilting

Types of Materials

Rock: hard or firm mass.

Debris: 20% to 80% of the particles are larger than 2mm, and the remainder is less than 2mm”.

Earth: material in which 80% or more of the particles are smaller than 2mm.

Soil: an aggregate of solid particles, generally of minerals and rocks.

Mud: material in which 80% or more of the particles are smaller than 0.06mm.

Types of flows

Debris flow: It is a form of rapid mass movement in which a combination of loose soil, rock, organic matter, slurry that flows downslope. They are commonly caused by intense precipitation or rapid snow melt.

Earth flow: It is down slope viscous flow of fine grained material saturated with water.

Mudflow: A mudflow is wet or viscous fluid mass of fine and coarse grained material flows rapidly along drainage channels.

Creep: Creep is the slow, steady, downward movement of material under gravity occurs in large area.

Lateral Spreads: It is the lateral movement usually occur on very gentle slopes or flat terrain. The failure is caused by liquefaction due to earthquake.



Figure 2. Landslides from different parts of India

RAPID RESPONSE TO LANDSLIDE DISASTER USING EO DATA




Figure 3. Landslide damage to New Haflong Railway Station Dima Hasao District of Assam (15-16 May 2022). Preliminary observation shows the occurrence of a number of landslides on the slopes on either side of the station. A large debris flow on the NW of the station has resulted in the overtopping and damage of a parked train in the station.




Figure 4. Damage caused by the Raunthi Gad landslide. Synoptic view of the affected region by Raunthi Gad rock fall and Debris flow (7 Feb 2021) as seen from Resourcesat-2A LISS IV image (08 February 2021) shown in FCC. Inset: shows enlarged view of the rock avalanche zone with joints and foliation.



Figure 5. Damage caused by the Raunthi Gad landslide a. Photograph taken from the helicopter showing the triangle shaped release area of the rock avalanche b. Foliation and joint-1 exposed on the joint-2 section, c. Ground photograph showing boulders of various sizes over the dam and on the valley floor, d. Tunnel filled with debris and water e. Ground photograph showing debris flow mark above the valley floor.




Figure 6. Geomorphological changes leading to the disaster around Kedarnath, Uttarakhand (2013). Satellite data reconstructed the chain of events around Kedarnath during 16-17 June 2013. It was found that reactivation of a large old landslide damaged a river training wall that resulted in the flooding of Kedarnath town on 16 June 2013. The twin events not only buried Kedarnath town with debris brought down from terminal and lateral moraines of Chorabari and Companion glaciers but also changed the course of Mandakini river from west of Kedarnath town to east of it. Pre-disaster Cartosat 1 image and post-disaster Cartosat-2S image of Kedarnath town showing structural damage to buildings and other facilities due to debris flow and flash flood.



Figure 7. Blockage of road (highlighted with circles) adjacent to the Tista river due to landslides triggered by Sikkim Earthquake, 2011 . Left - pre-earthquake WorldView-2 image, and right - post-earthquake pansharpened GeoEye-1 image shown with natural colour composite.




Figure 8. Parts of the road destroyed by landslides triggered due to the earthquake near Chungthang area in north Sikkim. Pre- and post-earthquake Cartosat-1 images are shown in left and right, respectively.




Figure 9. Landslide damage in Okhimath town. Pre- (left) and post- (right) landslide GeoEye-1 images showing buildings that were damaged (yellow outlines) by landslides. A. Mangali village - debris flow buried majority of houses, B. Chunni village - The debris flow that has buried Mangali village has also buried the majority of houses in Chunni village located in the downslope. C. Buildings in the debris flow path were levelled to the ground.

LANDSLIDE INVENTORY OF INDIA




Figure 10. Landslide inventory of India.

The database covers landslide-vulnerable regions of India in the Himalayas and Western Ghats. Satellite data of high to very high resolution nature such as IRS-1D PAN+LISS-III, Resourcesat-1, 2 and 2A LISS-IV Mx, Cartosat-1 and 2S, Aerial images were used in the mapping of landslides. Change detection using visual (manual) and digital (automatic) techniques were used to prepare the landslide inventory database. Some mapped landslides were validated in the field using mobile App and from the news report. The database mainly contains three types of inventory – seasonal, event-based and route-wise for the 2000-2017 period. Seasonal inventory contains pan-India landslide database corresponding to the 2014 and 2017 rainy season in India. The inventory database is available in a web GIS platform in the Bhuvan portal. The landslide inventory database shows the hotspot areas.

Table 2. Landslides inventory database of India.

Sl. No.	State/UTs	Monsoon season 2014	Monsoon season 2017	Field-based/year	Event-based / year	Total
1	Jammu and Kashmir	6826	19	434 / 2011	1 / 2015	7280
2	Ladakh	23		-	-	23
3	Himachal Pradesh	922	172	413 / 1998	1/ 2017 51/2013 2/2021	1561
4	Uttarakhand	1593	455	1419 / 1998	32/2003 307/2010 473 / 2012 6610 / 2013 1 / 2017 329/2021 1/2022	11219
5	Sikkim	73	79	-	1408 / 2011 8 / 2012 1/ 2016	1569
6	West Bengal	24	82	-	66 / 2011	172
7	Arunachal Pradesh	2904	4709	-	75 / 2016 1/2021	7689
8	Nagaland	54	2071	-	7/2017	2132
9	Manipur	379	4559	-	556/2017 1/2022	5494
10	Mizoram	1205	2254	-	8926/2017	12385
11	Tripura	56	8014	-	-	8070
12	Assam	1243	793	-	533/2017 5091/2022	2569
13	Meghalaya	2127	512	-	-	2639
14	Maharashtra	97	3	-	5012/2021	5112
15	Goa	2	1	-	-	3
16	Karnataka	82	19	-	993/2018	1094
17	Kerala	9	45	-	5191/2018 756/2019 09/2020 29/2021	6039
18	Tamil Nadu	79	8	-	603/2018	690
19	Haryana	-	100	-	-	100
	Total	17,698	23,895	2,266	37,074	80933




Figure 11. Landslides mapped using high-resolution satellite data in Jammu & Kashmir and Ladakh, which occurred between 2014 to 2017.



Figure 12. Paryote landslides near Paryote village in Doda district, Jammu and Kashmir. The continuous reactivation of Paryote Landslide near Paryote village in Doda district of Jammu and Kashmir. Translational-type failure with rock fall has occurred near the crown of the landslide, followed by a partially channelized debris flow along the main body of the landslide.




Figure 13. Landslides mapped using high-resolution satellite data in Himachal Pradesh, which occurred between 1998 to 2017.



Figure 14. Kotropi landslides, Mandi district, Himachal Pradesh. A massive landslide occurred near the village of Kotropi in Mandi district of Himachal Pradesh, on Sunday, 13th of August, 2017. The landslide occurred on National highway 154, the road between Mandi and Pathankot. The landslide resulted from rotation failure near the crown region followed by deep translational mechanism along the landslide body and channelised debris flow along the toe.




Figure 15. Landslides mapped using high-resolution satellite data in Uttarakhand, which occurred between 1998 to 2022.




Figure 16. Landslides around Kedarnath, Uttarakhand. During 16-17 June 2013, heavy rainfall (approximately 325 mm) caused widespread flash floods in the upstream areas of Mandakini, Alaknanda and Bhagirathi rivers resulting in massive damage to properties and infrastructures in the Kedarnath valley and areas in the downstream of Badrinath and Gangotri. A large reactivated landslide breached an artificial embankment north of the Kedarnath town.




Figure 17. Landslides mapped using high-resolution satellite data in Haryana, which occurred in 2017.




Figure 18. Panchkula landslides, Haryana. Landslide was triggered in the Panchkula of Haryana to heavy monsoonal rain in 2017. The landslide is shallow translational failure followed by channelized debris flow.




Figure 19. Landslides mapped using high-resolution satellite data in Sikkim, which occurred between 2011 to 2022.




Figure 20. Mantam landslide, Sikkim. A massive landslide occurred near the village of Mantam (opposite the Passingdang-Mantam road) in Sikkim on 13th of August, 2016. Wedge type failure has occurred near the crown of the landslide followed by a translational type of failure in the main body of the landslide.




Figure 21. Landslides mapped using high-resolution satellite data in West Bengal, which occurred between 2011 to 2017.




Figure 22. Landslide near Kurseong village, West Bengal. Reactivation of a large landslide near Kurseong town in Darjeeling district of West Bengal. Translational-type failure has occurred near the crown of the landslide, followed by a channelized debris flow along the main body of the landslide.




Figure 23. Landslides mapped using high-resolution satellite data in Arunachal Pradesh, which occurred between 2014 to 2021.




Figure 24. Landslide from West Siang, Arunachal Pradesh. Landslide was triggered in the West Siang of Arunachal Pradesh to heavy monsoonal rain in 2017. The landslide is shallow translational failure.




Figure 25. Landslides mapped using high-resolution satellite data in Assam, which occurred between 2014 to 2022.




Figure 26. Landslide from Dima Hasao District, Assam. Cluster landslides in Dima Hasao District of Assam caused by heavy rainfall in the month of May, 2022. Most of the landslides are shallow translational failures.




Figure 27. Landslides mapped using high-resolution satellite data in Nagaland, which occurred between 2014 to 2018.




Figure 28. Landslide from Kikruma from Phek district, Nagaland. A large landslide was triggered on 29 July 2018 near Kikruma village in Phek district of Nagaland resulting in the formation of an artificial dam on the Sidzu. The failure mechanism of the landslide is deep translational (planar) .




Figure 29. Landslides mapped using high resolution satellite data in Manipur which were occurred between 2014 to 2022.




Figure 30. Landslide Occurred near Tupul Railway Station, Noney, Manipur . A large landslide occurred near Tupul railway station in the Noney district of Manipur on 30 June 2022. From the geometry of the landslide, it can be inferred that the failure is a rotational type, and the crown is located on an unstable slope.




Figure 31. Landslides mapped using high-resolution satellite data in Mizoram, which occurred between 2014 to 2018.




Figure 32. Landslide from Aizawl, Mizoram. Landslide was triggered in the Aizawl of Mizoram to heavy monsoonal rain in 2017. The landslide is shallow translational failure.




Figure 33. Landslides mapped using high-resolution satellite data in Tripura which, occurred between 2014 to 2018.




Figure 34. Landslide from Dhalai, Tripura . Landslides were triggered in the Dhalai of Tripura to heavy monsoonal rain in 2017. The landslide is shallow translational failure.




Figure 35. Landslides mapped using high-resolution satellite data in Meghalaya, which occurred between 2014 to 2017.




Figure 36. Landslide from East Khasi Hills, Meghalaya. Landslides were triggered in the East Khasi Hills of Meghalaya due to heavy monsoonal rain in 2017. The landslides are shallow translational failures.




Figure 37. Landslides mapped using high-resolution satellite data in Maharashtra, which occurred between 2014 to 2021.




Figure 38. Landslide from Taliye Village ,Mahad district, Maharashtra. A catastrophic landslide occurred at the village of Taliye in Mahad taluk of Raigad on 22 July 2021. The landslide crown shows a rotational failure followed by deep translational failure along the main body.




Figure 39. Landslides mapped using high-resolution satellite data in Karnataka, which occurred between 2014 to 2018.




Figure 40. Landslide from Kodagu and Dakshina Kannada districts, Karnataka. Landslides were triggered in the Kodagu and Dakshina Kannada districts of Karnataka due to heavy rain in the 3rd week of August, 2018. Majority of the landslides were shallow translational failures.




Figure 41. Landslides mapped using high-resolution satellite data in Kerala which, occurred between 2014 to 2021.




Figure 42. Landslide from Puthumala region, Wayanad district, Kerala. Most of these landslides are channelised debris flows




Figure 43. Landslides mapped using high-resolution satellite data in Tamil Nadu, which occurred between 2014 to 2018.




Figure 44. Landslide in Coimbatore district, Tamil Nadu. Most of the landslides are shallow translational failures or channelised debris flows




Figure 45. Landslides mapped using high-resolution satellite data in Goa, which occurred between 2014 to 2017.

LANDSLIDE SUSCEPTIBILITY MAPPING AND DISTRICTS-WISE LANDSLIDE RISK EXPOSURE

Landslide Susceptibility Zonation

In view of the recurrence of landslides and resultant death toll in recent years in the Himalayas, and also appreciating the importance of remote sensing techniques in landslide related studies, at the instance of Cabinet Secretary, Government of India, Department of Space had undertaken a specific project for preparation of Landslide Susceptibility Zonation (LSZ) and Management maps on 1:25,000 scale in the Himalayas of Uttarakhand and Himachal Pradesh states.

The whole process of landslide phenomena is dependent on two groups of causative factors as below:

Geological / Topographic factors

- * Lithology
- * Geological structures (fault, trust, lineament etc)
- * Slope-dip (bedding, joint) relation
- * Geomorphology
- * Drainage
- * Slope angle, slope aspect and slope morphology
- * Land use / land cover
- * Soil texture and depth
- * Rock weathering

Triggering Factors

- * Anthropogeny
- * Rainfall
- * Earthquake

The geological factors are often considered for the Landslide Susceptibility Zonation, as they play a dominant role in the prognosis of landslide.

Keeping in view the various techniques of model, it has been felt that a model addressing the expert knowledge combined with a statistical technique suffices the need.

In this method, how each parameter of the terrain and categories within the parameters respond to landslide susceptibility are considered and ranked as per domain expert opinions. Finally, ranks are translated to weightages using the Decision space software and LSZ modeler for generation of the hazard maps. The basic principle used for developing this model is the Analytical Hierarchical Process (AHP) and Saaty's Principle of pair wise comparison.

Landslide Susceptibility Zonation

About 2000 km stretch of routes was taken up for LSZ study. For convenience in handling the huge database, the routes were divided into different sectors

Uttarakhand

1. Rishikesh – Rudraprayag – Chamoli – Badrinath
2. Rudraprayag – Okhimath – Kedarnath
3. Chamoli – Okhimath
4. Rishikesh – Uttarkashi – Gangotri – Gaumukh
5. Pithoragarh – Khela - Malpa

Himachal Pradesh

1. Shimla – Rampur – Sarahan – Sumdo
2. Shimla – Bilaspur – Kulu – Manali
3. Dalhousie – Chamba – Brahmaur




Figure 46. LSZ map for Shimla – Rampur – Sarahan – Sumdo Route.

Landslide risk exposure

The occurrence of landslides in an area is due to the interplay of favorable terrain parameters such as slope, lithology, topography and land use which trigger landslides in response to rainfall or earthquake events.

Preparation of landslide inventory, hazard, and risk mapping is done in India using satellite data to analyze the cause of landslides which is mostly found to be :

- Litho-tectonic control- earthquake
- Precipitation (rainfall) control.

In India, the major areas which are affected by landslides are :

- The Northwest Himalayas contribute- 66.5% of landslides in India, followed by the Northeast Himalayas -18.8% and the Western Ghats - 14.7%.

Methods of mapping landslide inventory

Data collected: pre-monsoon season (Mar–May, 2014) and post-monsoon season (Oct–Dec, 2014).

Satellites used: Resourcesat-2 LISS-IV Mx (5.8 m)- for seasonal landslide mapping - trigger by monsoon rainfall.

As manual method of landslide mapping is a tedious job, so semi-automatic method developed by Martha et al. (2010, 2011, 2012) for the detection of landslides from high-resolution satellite data and DEM was used for the creation of the seasonal landslide inventory database.

Event-based landslide (caused by earthquakes or short-duration rainfall event) inventory mapping:

An event-based landslide inventory map was prepared using the same method as adopted for the SLIM.

Field-based landslide inventory:

This landslide inventory tallies to 1998, which was prepared by NRSC by means of merged IRS-1D PAN+LISS III data (5.8 m) by visual image interpretation and verified on the ground (NRSA 2001).

Assessment of Socio-economic Parameter exposure

Socio-economic parameters (SEP) include total population, number of households, road and livestock. The unit of exposure analysis is taken as district and two indices were created so that they normalize with the area of the same district. These Indices are :

$$1) \text{ District landslide density (DLD)} = \frac{\text{Area of landslide in district}}{\text{Area of district}}$$

$$2) \text{ District SEP density (DSEPD)} = \frac{\text{SEP count}}{\text{Area of district}}$$

These indices were calculated using excel table operation.

Later district exposure index (DEI) was calculated to rank the districts.

$$\text{District Exposure Index} = \text{DLD} * (\text{DSEPD}_p + \text{DSEPD}_h + \text{DSEPD}_l + \text{DSEPD}_r)$$

where subscripts p, h, l and r in DSEPD refer to population, house, livestock and road, respectively.

Landslide risk exposure




Figure 47. Socio-economic parameter risk exposure map, The contribution of each exposure element for the top ten landslide exposed districts is shown in bar diagram.

Socio-economic parameter exposure

Landslide exposure analysis were carried out in the mountainous areas. Rudraprayag district in Uttarakhand state which has highest landslide density in India is also having highest exposure to total population, working population, literacy and no. of houses. Below is the rank of all 147 districts in 17 states and 02 UTs of India for their exposure to landslides in terms of key socio economic parameters.

Table 3. Table shows the rank of districts of India for their exposure to landslides.

District Rank	District	State	District Rank	District	State
1	Rudraprayag	Uttaranchal	36	Coimbatore	Tamil Nadu
2	Tehri Garhwal	Uttaranchal	37	Solan	Himachal Pradesh
3	Thrissur	Kerala	38	Aizawl	Mizoram
4	Rajauri	J&K	39	Lunglei	Mizoram
5	Palakkad	Kerala	40	Kamrup	Assam
6	Poonch	J&K	41	Dindigul	Tamil Nadu
7	Malappuram	Kerala	42	Kathua	J&K
8	South District	Sikkim	43	Kanniyakumari	Tamil Nadu
9	East District	Sikkim	44	Kasaragod	Kerala
10	Kozhikode	Kerala	45	Lawngtlai	Mizoram
11	Imphal West	Manipur	46	Kinnaur	Himachal Pradesh
12	Kodagu	Karnataka	47	Hailakandi	Assam
13	Wayanad	Kerala	48	Kollam	Kerala
14	Jammu	J&K	49	Goalpara	Assam
15	Ernakulam	Kerala	50	Bageshwar	Uttaranchal
16	Mandi	Himachal Pradesh	51	North District	Sikkim
17	Udhampur	J&K	52	Anantanag	J&K
18	Idukki	Kerala	53	Hassan	Karnataka
19	Chamoli	Uttaranchal	54	Dakshina Kannada	Karnataka
20	West District	Sikkim	55	Karbi Anglong	Assam
21	Uttarkashi	Uttaranchal	56	Lohit	Arunachal Pradesh
22	Cachar	Assam	57	Kullu	Himachal Pradesh
23	Garhwal	Uttaranchal	58	Baramulla	J&K
24	Kottayam	Kerala	59	Theni	Tamil Nadu
25	Hamirpur	Himachal Pradesh	60	Kolasib	Mizoram
26	Kannur	Kerala	61	Shimla	Himachal Pradesh
27	Pulwama	J&K	62	Kangra	Himachal Pradesh
28	Thiruvananthapuram	Kerala	63	Churachandpur	Manipur
29	Dehra Dun	Uttaranchal	64	East Garo Hills	Meghalaya
30	Bilaspur	Himachal Pradesh	65	Champawat	Uttaranchal
31	West Garo Hills	Meghalaya	66	West Khasi Hills	Meghalaya
32	Chamba	Himachal Pradesh	67	Ribhoi	Meghalaya
33	Pathanamthitta	Kerala	68	Naini Tal	Uttaranchal
34	East Khasi Hills	Meghalaya	69	Jaintia Hills	Meghalaya
35	Darjiling	West Bengal	70	Una	Himachal Pradesh

Table 3. Table shows the rank of districts of India for their exposure to landslides continues.....

District Rank	District	State	District Rank	District	State
71	Mamit	Mizoram	110	Dhalai	Tripura
72	Tirunelveli	Tamil Nadu	111	North Goa	Goa
73	Papum Pare	Arunachal Pradesh	112	Ukhrul	Manipur
74	Tawang	Arunachal Pradesh	113	Kurung Kumey	Arunachal Pradesh
75	West Tripura	Tripura	114	Sindhudurg	Maharashtra
76	North Tripura	Tripura	115	Tirap	Arunachal Pradesh
77	Udupi	Karnataka	116	Uttara Kannada	Karnatka
78	East Siang	Arunachal Pradesh	117	Nagaon	Assam
79	Doda	J&K	118	Champhai	Mizoram
80	Thane	Maharashtra	119	Budgam	J&K
81	Almora	Uttaranchal	120	Upper Subansiri	Arunachal Pradesh
82	Serchhip	Mizoram	121	South Goa	Goa
83	North Cachar Hills	Assam	122	Chandel	Manipur
84	Lower Subansiri	Arunachal Pradesh	123	Kargil	J&K
85	The Nilgiris	Tamil Nadu	124	Haveri	Karnataka
86	Pithoragarh	Uttaranchal	125	Anjaw	Arunachal Pradesh
87	West Kameng	Arunachal Pradesh	126	Lahul & Spiti	Himachal Pradesh
88	Sirmaur	Himachal Pradesh	127	Dhubri	Assam
89	Phek	Nagaland	128	Nashik	Maharashtra
90	South Garo Hills	Meghalaya	129	Ratnagiri	Maharashtra
91	Changlang	Arunachal Pradesh	130	South Tripura	Tripura
92	Chikmagalur	Karnataka	131	Ahmadnagar	Maharashtra
93	Mon	Nagaland	132	Kupwara	J&k
94	Tuensang	Nagaland	133	Kolhapur	Maharashtra
95	Bongaigaon	Assam	134	Satara	Maharashtra
96	Lower Dibang Valley	Arunachal Pradesh	135	Dibang Valley	Arunachal Pradesh
97	Senapati	Manipur	136	Leh-ladakh	J&K
98	Srinagar	J&k	137	Karimganj	Assam
99	Morigaon	Assam	138	Alappuzha	Kerala
100	Tamenglong	Manipur	139	Mumbai Suburban	Maharashtra
101	East Kameng	Arunachal Pradesh	140	Mumbai	Maharashtra
102	Saiha	Mizoram	141	Thoubal	Manipur
103	Shimoga	Karnataka	142	Imphal East	Manipur
104	Upper Siang	Arunachal Pradesh	143	Bishnupur	Manipur
105	Mokokchung	Nagaland	144	Zunheboto	Nagaland
106	Kohima	Nagaland	145	Wokha	Nagaland
107	Pune	Maharashtra	146	Haridwar	Uttaranchal
108	West Siang	Arunachal Pradesh	147	Udham Singh Nagar	Uttaranchal
109	Raygarh	Maharashtra			

RESEARCH & DEVELOPMENT IN LANDSLIDES STUDIES

Multitemporal SAR interferometry for Landslide prediction

Landslides from remote steep slopes render people living downhill vulnerable, unaware of the impending danger. Identifications of slow-moving mountain slopes are possible now due to time series measurements from space using microwave satellite data and the InSAR technique, which can detect displacement at the millimetre level. The availability of open-source Sentinel-1 data has revolutionised the study involving landslide kinematics and predicting the time of failure. However, identifying accelerating trends, demarcation of the release area, and predicting flow path after failure initiation is still challenging.

Time of failure prediction of the Kikrma landslides in the NE Himalaya is carried out using Persistent Scatter Interferometry (PSI) interferometric technique on Sentinel-1 SAR data to analyse the trend of ground deformation leading to slope failures. The displacement time series of the measurement points (MPs), analysed using inverse velocity and modified inverse velocity methods, show that the instability had commenced almost a year or more with the final onset of acceleration (OOA) triggered by heavy rainfall, couple of weeks prior to the actual failure (Roy et al. 2022).




Figure 48. Landslide prediction using MT InSAR. Top panel: Representative displacement time series and inverse velocities for the Kikrma landslide showing the OOA (t_0) and last observation point (t). Inset: Failure day estimation using INV. Bottom panel: Rainfall variation in the region during the observation period. Predicted day of failure: 26 July 2018; Actual day of failure: 29 July 2018.

Debris Flow Modelling

Dynamic modelling of debris flow landslides has become an increasingly important practice for simulating the characteristics and behaviour of debris flow relying on the physical laws of conservation of mass, momentum and energy (Melo et al., 2018). This model approach can be broadly classified as the lumped mass moment where motion of slide is considered as a single point which is not always the case or based on continuum mechanics associated with a distinct rheology (Hung, 1995). To simulate the deformation of the moving mass along the flow path, choosing appropriate friction parameters and material rheologies is taxing (Rickenmann, 2005).

Numerical debris flow modelling was carried out for several major landslides in Kerala to characterize the landslide geometry and assess their impact.

For modeling debris flow paths, Rapid Mass Movement Simulation (RAMMS) software developed by WSL Institute for Snow and Avalanche Research SLF and the Swiss Federal Institute for Forest, Snow and Landscape Research WSL, was used. The model gives flow height, velocity and pressure of debris flow. The main input parameters to be ingested in the model are high resolution Digital Elevation Model (DEM) for deriving topographic parameters, release area and friction information. CartoDEM (2.5 m) was used for flow modeling (Jain et al. 2021).




Figure 49. Flow modeling for Puthumala Landslide, Wayanad District, Kerala.

UAV data in landslide modeling

Photogrammetric techniques have been increasingly used because of their capability to rapidly reconstruct the 3-D topography from aerial photographs and provided such data exist for different time periods, allow objective change detection (Martha et al, 2010). Knowledge of failure volumes is also critical for a more accurate understanding of the landslide process and its flow, which shows the affected areas.

The increasing availability of high-resolution UAV data, quantitative studies on landform changes using DEMs have become a viable option.

UAV data were acquired over the Puthumala landslide in Kerala area on 22 March 2021. The resolution of orthoimage of UAV is 2 cm and the grid size of DTM is 8 cm. The positional accuracy of the image is 1-2 m. . Pre and post-DEM were utilized for volumetric estimation and release area mapping.




Figure 50. Landslide modeling using UAV data, Puthumala Landslide, Wayanad District, Kerala.

Landslide Susceptibility Zonation: New Approach

A large area covering 21,087 km² of the Mizoram state, India has been investigated, using five statistical methods of susceptibility models viz. Multi Criterion Weighted Overlay (MCWO), Information Value (IV), Weights of Evidence (WoE), Logistic Regression (LR) and Artificial Neural Network (ANN) to prepare macro-scale landslide susceptibility maps





Figure 51. Landslide susceptibility maps generated using spatial sampling strategy by a) MCWO, b) IV, c) WoE, d) LR and e) ANN. f - prediction and success rate curves for of all five models.

Rainfall-Threshold based landslide early warning

A regional landslide early warning system for selective routes of HP, Uttarakhand and NER is operational on an experimental basis for monsoon season. It uses rainfall forecast data from MOSADAC, IMD and Climate Prediction Center (CPC). Below is the landslide early warning was given along the Dalhousie-Chamba route.

Rainfall threshold for landslides is the value which when reached or exceeded is likely to trigger landslides. Rainfall thresholding for slope failure can be established using process based, empirical or statistical approach. Historical data on landslide causing rainfall events and the corresponding landslide records can be used to establish rainfall thresholds for various geographical extents based on consideration of specific rainfall events or using antecedent conditions. The I-D based thresholding is the most widely used empirical thresholding method. The thresholding is done by drawing lower-bound lines on plots of points representing landslide triggering rainfall events. The daily, 3-day cumulative and 15-day & 30-day antecedent rainfall values associated with landslides had been subjected to binary logistic regression using landslide as dichotomous dependent variable. significant predictors influencing slope failure. By using binary logistic regression model the slope failure triggering probability is estimated.



The logistic regression retained the daily (DR), 3-day cumulative (3DCR) and 30-day antecedent rainfall (30DAR) as significant predictors influencing slope failure.

$$z = -3.817 + DR * 0.077 + 3DCR * 0.058 + 30DAR * 0.009$$

$$f(z) = \frac{1}{1 + e^{-z}}, \quad z: -\infty \text{ to } +\infty, P: 0 \text{ to } 1$$

Different combinations of triggering probability and the Landslide Susceptibility classes used in this approach are described below:

Figure 52. Landslide early warning for advance 3 days given in parts of Himachal Pradesh.



Landslide in Shimla

Landslide Susceptibility	Triggering Probability	
	0.75 to 0.85	> 0.85
Severe	Warn	Warn
Very High	Watch	Warn
High	Watch	Watch
Moderate	Advisory	Advisory

Table 4. Table showing triggering probability for landslide trigger.

Mobile Application for Landslide data collection



Figure 53. Step-by-step procedure for landslide data collection using FLIM mobile app.

The FLIM mobile app is developed by NRSC/ISRO. It aimed to be used by SDMAs/DDMAs/Govt. organizations/researchers with reasonable knowledge on landslides. The main purposes of this app are to collect data of landslides from the field and to develop a landslide warning system for the region using the collected field landslide data.

Mobile Application for Landslide data collection



Figure 54. Landslide data from western Maharashtra collected in Bhuvan FLIM database.



Figure 55 . Ground landslide data collected in NER using FLIM mobile App.

The FLIM mobile app is developed by NRSC/ISRO. It aimed to be used by SDMAs/DDMAs/Govt. organizations/researchers with reasonable knowledge on landslides. The main purposes of this app are to collect data of landslides from the field and to develop a landslide warning system for the region using the collected field landslide data. The collected data sent from the field using this app will be uploaded in the BHUVAN and NDEM servers for analysis and visualization. Hence, the app will help the state agencies as incident reporting system for landslide disasters and for development of a landslide early warning system in the region.

References

- D. M. Cruden. A Simple Definition of a Landslide. *Bulletin of the International Association of Engineering Geology*, Volume 43, 1991, 27-29. <https://doi.org/10.1007/BF02590167>.
- D. J. Varnes. Slope movement types and processes, In: R.L. Schuster and R.J. Krizek, *Landslides—Analysis and control*. National Research Council, Special Report 176, 1978, 11–33.
- D. J. Varnes. *Landslide Types and Processes, Landslides and Engineering Practice*. National Research Council, Special Report 58, 1958, 20-47.
- D. Rickenmann. Runout prediction methods. In: *Debris-flow Hazards and Related Phenomena*. Springer Praxis Books. Springer, Berlin, Heidelberg, 2005. https://doi.org/10.1007/3-540-27129-5_13.
- Fabio Vittorio De Blasio. *Introduction to the Physics of Landslides, Lecture Notes on the Dynamics of Mass Wasting*, Springer Dordrecht Heidelberg London New York. 2011. <https://doi.org/10.1007/978-94-007-1122-8>.
- F. Nadim, O. Kjekstad, P. Peduzzi, C. Herold, C. Jaedicke. Global landslide and avalanche hotspots. *Landslides* 3, 2006, 159–173. <https://doi.org/10.1007/s10346-006-0036-1>.
- L. M. Highland, P. Bobrowsky. *The Landslide Handbook - A Guide to Understanding Landslides*. U.S. Geological Survey, 2008.
- Nirmala Jain, Tapas Ranjan Martha, Kriti Khanna, Priyom Roy, K. Vinod Kumar. Major landslides in Kerala, India, during 2018–2020 period: an analysis using rainfall data and debris flow model. *Landslides* 18, 2021, 3629–3645. <https://doi.org/10.1007/s10346-021-01746-x>.
- Oldrich Hungr. A Model for the Runout Analysis of Rapid Flow Slides, Debris Flow, and Avalanches. *Canadian Geotechnical Journal*, 1995, Volume 32, 610-623. <https://doi.org/10.1139/t95-063>.
- Oldrich Hungr, Serge Leroueil, Luciano Picarelli. The Varnes classification of landslide types, an update. *Landslides* 11, 2014, 167–194. <https://doi.org/10.1007/s10346-013-0436-y>.
- Oldrich Hungr, S. G. Evans, M. J. Bovis, J. N. Hutchinson. A review of the classification of landslides of the flow type. *Environmental & Engineering Geoscience*, Volume 7, 2001, 221–238. <https://doi.org/10.2113/gseegeosci.7.3.221>.

Pierluigi Confuorto. 'From site-scale to large areas monitoring of ground deformation phenomena by integration of different DInSAR techniques in Crotona Province (Southern Italy)'. 2016. <https://doi.org/10.13140/RG.2.1.2146.1364>.

Tapas Ranjan Martha, Priyom Roy, Nirmala Jain, Kirti Khanna, K. Mrinalni, K. Vinod Kumar ,P. V. N. Rao. Geospatial landslide inventory of India—an insight into occurrence and exposure on a national scale. *Landslides* 18, 2021, 2125–2141. <https://doi.org/10.1007/s10346-021-01645-1>.

Tapas Ranjan Martha, Norman Kerle, Victor Jetten, Cees J. van Westen, K. Vinod Kumar. Characterising spectral, spatial and morphometric properties of landslides for semi-automatic detection using object-oriented methods. *Geomorphology*, Volume 116, 2010, 24-36. <https://doi.org/10.1016/j.geomorph.2009.10.004>.

Tapas Ranjan Martha, Norman Kerle, Cees J. van Westen, Victor Jetten, K. Vinod Kumar. Object-oriented analysis of multi-temporal panchromatic images for creation of historical landslide inventories, *ISPRS Journal of Photogrammetry and Remote Sensing*, Volume 67, 2012, 105-119, <https://doi.org/10.1016/j.isprsjprs.2011.11.004>.

Turner, A. Keith, and Schuster, L. Robert. *Landslides—Investigation and mitigation*. National Research Council, National Academy Press, 1996.

List of Websites

National Remote Sensing Centre (NRSC) portal: <https://www.nrsc.gov.in/>

Bhuvan portal: <https://bhuvan.nrsc.gov.in/home/index.php>

National Database for Disaster Management (NDEM): <https://ndem.nrsc.gov.in/>

Bibliography of landslide studies done by NRSC

- Priyom Roy, Tapas Ranjan Martha, K. Vinod Kumar, Prakash Chauhan, Vala Venkateshwar Rao. Cluster landslides and associated damage in the Dima Hasao district of Assam, India due to heavy rainfall in May 2022. *Landslides* 20, 2023, 97–109. <https://doi.org/10.1007/s10346-022-01977-6>.
- Priyom Roy, Tapas Ranjan Martha, Kirti Khanna, Nirmala Jain, K. Vinod Kumar. Time and path prediction of landslides using InSAR and flow model, *Remote Sensing of Environment*. Volume 271, 2022, 112899. <https://doi.org/10.1016/j.rse.2022.112899>.
- Priyom Roy, Nirmala Jain, Tapas Ranjan Martha, K. Vinod Kumar. Reactivating Balia Nala landslide, Nainital, India—a disaster in waiting. *Landslides* 19, 2022, 1531–1535. <https://doi.org/10.1007/s10346-022-01881-z>.
- Lina Hao, Cees van Westen, A. Rajaneesh, K.S. Sajinkumar, Tapas Ranjan Martha, Pankaj Jaiswal. Evaluating the relation between land use changes and the 2018 landslide disaster in Kerala, India. *CATENA*, Volume 216, 2022, 106363. <https://doi.org/10.1016/j.catena.2022.106363>.
- Vipin Kumar Maurya, Ramji Dwivedi, Tapas Ranjan Martha. Site scale landslide deformation and strain analysis using MT-InSAR and GNSS approach – A case study. *Advances in Space Research*. Volume 70, 2022, 3932-3947. <https://doi.org/10.1016/j.asr.2022.05.028>.
- S. Devaraj, K. Yarrakula, Tapas Ranjan Martha, Geetha Priya Murugesan, Divya Sekhar Vaka, Samvedya Surampudi, Abhinav Wadhwa, Parthiban Loganathan, Venkatesh Budamala. Time series SAR interferometry approach for landslide identification in mountainous areas of Western Ghats, India. *J Earth Syst Sci* 131, 2022, 133. <https://doi.org/10.1007/s12040-022-01876-3>.
- Tapas Ranjan Martha, Priyom Roy, Nirmala Jain, Kirti Khanna, K. Mrinalni, K. Vinod Kumar, P. V. N. Rao. Geospatial landslide inventory of India—an insight into occurrence and exposure on a national scale. *Landslides* 18, 2021, 2125–2141. <https://doi.org/10.1007/s10346-021-01645-1>.
- Tapas Ranjan Martha, Priyom Roy, Nirmala Jain, K. Vinod Kumar, P. Sashivardhan Reddy, J. Nalini, S. V. S. P. Sharma, Abhinav Kumar Shukla, K. H. V. Durga Rao, B. Narender, P. V. N. Rao, S. Muralikrishnan. Rock avalanche induced flash flood on 07 February 2021 in Uttarakhand, India—a photogeological reconstruction of the event. *Landslides* 18, 2021, 2881–2893. <https://doi.org/10.1007/s10346-021-01691-9>.

- Nirmala Jain, Tapas Ranjan Martha, Kriti Khanna, K. Vinod Kumar. Major landslides in Kerala, India, during 2018–2020 period: an analysis using rainfall data and debris flow model. *Landslides* 18, 2021, 3629–3645. <https://doi.org/10.1007/s10346-021-01746-x>.
- Alessandro Cesare Mondini, Fausto Guzzetti, Kang-Tsung Chang, Oriol Monserrat, Tapas Ranjan Martha, Andrea Manconi. Landslide failures detection and mapping using Synthetic Aperture Radar: Past, present and future. *Earth-Science Reviews*. Volume 216, 2021, 103574. <https://doi.org/10.1016/j.earscirev.2021.103574>.
- Kriti Khanna, Tapas Ranjan Martha, Priyom Roy, K. Vinod Kumar. Effect of time and space partitioning strategies of samples on regional landslide susceptibility modeling. *Landslides* 18, 2021, 2281–2294. <https://doi.org/10.1007/s10346-021-01627-3>.
- Lina Hao, A. Rajaneesh, Cees van Westen, K.S Sajinkumar, Tapas Ranjan Martha, Pankaj Jaiswal, and Brian G. McAdoo. Constructing a complete landslide inventory dataset for the 2018 monsoon disaster in Kerala , India, for land use change analysis. *Earth System Science Data*, Volume 12, 2020, 2899–2918. <https://doi.org/10.5194/essd-12-2899-2020>.
- Tapas Ranjan Martha, Priyom Roy, Kirti Khanna, K. Mrinalni and K. Vinod Kumar. “Landslides Mapped Using Satellite Data in the Western Ghats of India after Excess Rainfall during August 2018.” *Current Science*, Volume 117, 2019, 804–12. URL: <https://www.jstor.org/stable/27138344>.
- Ren N. Keyport, Thomas Oommen, Tapas Ranjan Martha, K.S. Sajinkumar, John S. Gierke. A comparative analysis of pixel and object-based detection of landslides from very high-resolution images. *International Journal of Applied Earth Observation and Geoinformation*, Volume 64, 2018, 1-11, <https://doi.org/10.1016/j.jag.2017.08.015>.
- Priyom Roy, Tapas Ranjan Martha, Nirmala Jain and K. Vinod Kumar. “Reactivation of Minor Scars to Major Landslides – a Satellite-Based Analysis of Kotropi Landslide (13 August 2017) in Himachal Pradesh, India.”. *Current Science*, Volume 115, 2018, 395–98. URL: <https://www.jstor.org/stable/26978223>.
- Tapas Ranjan Martha, Priyom Roy, Ritwik Mazumdar, K. Babu Govindharaj, K. Vinod Kumar. Spatial characteristics of landslides triggered by the 2015 Mw 7.8 (Gorkha) and Mw 7.3 (Dolakha) earthquakes in Nepal. *Landslides* 14, 2017, 697–704. <https://doi.org/10.1007/s10346-016-0763-x>.

- Tapas Ranjan Martha, P. Shashivardhan Reddy, C. M. Bhatt, K. Babu Govindha Raj, J. Nalini, E. Anantha Padmanabha, B. Narender, K. Vinod Kumar, S. Muralikrishnan, G. Srinivasa Rao, P. G. Diwakar, V. K. Dadhwal. Debris volume estimation and monitoring of Phuktal river landslide-dammed lake in the Zaskar Himalayas, India using Cartosat-2 images. *Landslides* 14, 2017, 373–383. <https://doi.org/10.1007/s10346-016-0749-8>.
- Tapas Ranjan Martha, P. Kamala, Josna Jose, K. Vinod Kumar, G. Jai Sankar. Identification of new Landslides from High Resolution Satellite Data Covering a Large Area Using Object-Based Change Detection Methods. *J Indian Soc Remote Sens* 44, 2016, 515–524. <https://doi.org/10.1007/s12524-015-0532-7>.
- Tapas Ranjan Martha, Priyom Roy, K. Babu Govindharaj, K. Vinod Kumar, P. G. Diwakar, V. K. Dadhwal . Landslides triggered by the June 2013 extreme rainfall event in parts of Uttarakhand state, India. *Landslides* 12, 2015, 135–146 . <https://doi.org/10.1007/s10346-014-0540-7>.
- Tapas Martha Ranjan, K. Vinod Kumar. September, 2012 landslide events in Okhimath, India—an assessment of landslide consequences using very high resolution satellite data. *Landslides* 10, 2013, 469–479. <https://doi.org/10.1007/s10346-013-0420-6>.
- Tapas Ranjan Martha, Cees J. van Westen, Norman Kerle, Victor Jetten, K. Vinod Kumar. Landslide hazard and risk assessment using semi-automatically created landslide inventories. *Geomorphology*, Volume 184, 2013, 139-150. <https://doi.org/10.1016/j.geomorph.2012.12.001>.
- Tapas Ranjan Martha, Cees J. van Westen, Norman Kerle, Victor Jetten, K. Vinod Kumar. Landslide hazard and risk assessment using semi-automatically created landslide inventories. *Geomorphology*, Volume 184, 2013, 139-150. <https://doi.org/10.1016/j.geomorph.2012.12.001>.
- Shraban Sarkar, Archana K. Roy, Tapas Ranjan Martha. Landslide susceptibility assessment using Information Value Method in parts of the Darjeeling Himalayas. *J Geol Soc India* 82, 2013, 351–362 . <https://doi.org/10.1007/s12594-013-0162-z>.
- Tapas Ranjan Martha, Norman Kerle, Cees J. van Westen, Victor Jetten, K. Vinod Kumar. Object-oriented analysis of multi-temporal panchromatic images for creation of historical landslide inventories, *ISPRS Journal of Photogrammetry and Remote Sensing*, Volume 67, 2012, 105-119, <https://doi.org/10.1016/j.isprsjprs.2011.11.004>.

Tapas Ranjan Martha, Norman Kerle, Cees J. van Westen, Victor Jetten, K. Vinod Kumar. Segment Optimization and Data-Driven Thresholding for Knowledge-Based Landslide Detection by Object-Based Image Analysis. *IEEE Transactions on Geoscience and Remote Sensing*, Volume 49, 2011, 4928-4943. <https://doi.org/10.1109/TGRS.2011.2151866>.


Tapas Ranjan Martha, Norman Kerle, Victor Jetten, Cees J. van Westen, K. Vinod Kumar. Characterising spectral, spatial and morphometric properties of landslides for semi-automatic detection using object-oriented methods. *Geomorphology*, Volume 116, 2010, 24-36. <https://doi.org/10.1016/j.geomorph.2009.10.004>.

Tapas Ranjan Martha, Norman Kerle, Victor Jetten, Cees J. van Westen, K. Vinod Kumar. Landslide Volumetric Analysis Using Cartosat-1-Derived DEMs. *IEEE Geoscience and Remote Sensing Letters*, Volume 7, 2010, 582-586. <https://doi.org/10.1109/LGRS.2010.2041895>.

K. Vinod Kumar, R. C. Lakhera, Tapas Ranjan Martha, R. S. Chatterjee, A. Bhattacharya . Analysis of the 2003 Varunawat Landslide, Uttarkashi, India using Earth Observation data. *Environ Geol* 55, 2008, 789–799 . <https://doi.org/10.1007/s00254-007-1032-z>.



Landslide near Jakholi village, near Agastyamuni town, Rudraprayag district, Uttarakhand




Landslide from Bhiri village, near Guptakashi, Mandakini Valley, Uttarakhand




Landslide is located in Mohand ridge in Sivallik hills near Deharadun, Uttarakhand



Landslide from Chamoli district, Uttarakhand




Landslide near Jakholi village, near Agastyamuni town, Rudraprayag district, Uttarakhand



Landslide from Bhiri village, near Guptakashi, Mandakini Valley, Uttarakhand



Landslide is located in Mohand ridge in Sivalik hills near Deharadun, Uttarakhand



Landslide from Chamoli district, Uttarakhand